

Distribution of Cold dust in Orion A and B

B. Mookerjee¹, S.K. Ghosh¹, T.N. Rengarajan¹,
S.N. Tandon² & R.P. Verma¹

ABSTRACT

Large scale far-infrared (FIR) observations of the Orion complex at 205 and 138 μm are presented with an aim of studying the distribution of cold (<25 K) dust. The maps in these FIR bands extend over ~ 3600 sq. arcmin and cover regions around OMC-1, 2, 3 in Orion A and NGC 2023 and NGC 2024 in Orion B. Some limited regions have also been mapped at 57 μm . A total of 15 sources in Orion A and 14 in Orion B (south) have been identified from our FIR maps. Dust temperature distribution in both Orion A and Orion B (south) have been determined reliably using the maps at 205 & 138 μm obtained from simultaneous observations using almost identical beams (1.6 dia). These temperatures have been used to generate map of τ_{150} , the optical depth at 150 μm , for the Orion B region. The coldest source detected is in OMC-3 and has a temperature of ~ 15 K. The diffuse FIR emission in the different sub-regions is found to vary between 25 % to 50 % of the total FIR emission from that sub-region.

Subject headings: infrared: ISM: continuum – ISM: individual (OMC-1, OMC-2, OMC-3, NGC 2024, NGC 2023)

1. Introduction

In this paper we present a large scale high angular resolution ($\sim 1'$) study of the continuum emission at 205 and 138 μm from the two Giant Molecular Clouds Orion A and Orion B (south) in the Orion complex. The mapped region of Orion A includes OMC-1, 2, 3 and IRAS 05327-0457 (NGC 1977). In Orion B we have mapped mainly the southern portion covering NGC 2023 and NGC 2024. Limited regions including, OMC-1 and 2 in Orion A and NGC 2024 in Orion B have also been observed at 57 μm .

The southern part of the Orion cloud complex, also known as Orion A, contains star forming cores which are distributed along the north-south direction (integral shaped ridge) and named OMC-1, 2 and 3 from south to north. Observations at sub-millimeter/far-infrared wavelengths have shown that both warm and cold dust with quite different spatial distribution contribute to the emission from the Orion molecular clouds (Thronson et al. 1986; Chini et al. 1997). Cold dust, with temperature ~ 20 -25 K, accounts for bulk of the mass

of dust in the Orion clouds and emits significantly only at wavelengths longer than 100 μm . Earlier observations of Orion A concentrated on the emission from the hot dust viz., mapping of OMC-1 at 50-100 μm by Werner et al. (1976) and of M43 at 37-160 μm by Smith, Harper, & Loewenstein (1987). Later observations have mostly explored the emission from the cold dust in this region. The most recent of these observations was by Johnstone & Bally (1999) covering OMC-1, 2 and 3 and extending over an area of $50' \times 10'$ with an angular resolution of $\sim 7''.5$ at 450 μm and $14''$ at 850 μm . Lis et al. (1998) had also mapped a region of $6' \times 18'$ covering OMC-2 and 3 at 350 μm with an angular resolution of $11''$. However, the pioneering work in detecting a number of cold sources in OMC-2 and an extremely cold one (~ 10 K) in OMC-3 was by Chini et al. (1997), who mapped a region of $\sim 15' \times 5'$ with an angular resolution of $\sim 11''$ at 1300 μm . Most of these condensations are considered to be protostellar candidates because of the associated outflow activities (Castets & Langer 1995). Observations in molecular lines like C^{18}O by (Castets & Langer 1995, and references therein) have revealed that the gas in OMC-2 and OMC-3 have kinetic temperatures equal to 24 K and 19 K respectively, both much smaller than that in OMC-1 (70 K). These observations also identified

¹Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India

²Inter-University Centre for Astronomy & Astrophysics, Ganeshkhind, Pune 411 007, India

both, OMC-2 and OMC-3, to be regions of high column density. $C^{18}O$ observations have also detected several moderate to strongly collimated bipolar outflows in OMC-3, which suggests recent star formation activity. The very low temperatures of sources in both OMC-2 and 3 along with the evidence for star formation, indicate very early stages of evolution of, probably intermediate mass stars. In this paper we have explored the distribution of dust emission and temperature of the Orion A region in a wavelength regime (205 & 138 μm in particular) where no high angular resolution measurements exist in literature.

The Orion B cloud contains 4 major sites of star formation viz., NGC 2071 & NGC 2068 in the north and NGC 2023 & NGC 2024 in the south. We have observed only the regions around NGC 2024 and 2023. The brightest emission source in the region is NGC 2024, which is a region of active star formation. The 1300 μm observations over a region of $\sim 5' \times 5'$ (with an angular resolution $\sim 30''$) by Mezger et al. (1988) have revealed six condensations in NGC 2024, aligned in the north-south direction. These condensations also have masers and HII regions associated with them. The nature of these condensations was an issue of debate which was resolved by Mezger et al. (1992). From their observations at 870 & 1300 μm they concluded that these condensations are indeed isothermal protostars. Lada, Bally, & Stark (1991a) presented a survey of Orion B (Lynds 1630) molecular cloud in CS(2-1) line (1.8 angular resolution) and identified the regions including NGC 2023 and NGC 2024 to be bright in CS, thus indicating high density of the region. In a follow-up program Lada et al. (1991b) surveyed Orion B in the K (2.2 μm) band and found that more than 50% of the NIR sources are associated with the 4 identified star forming regions. In a recent high angular resolution (22-30'') survey of Orion B (south) at 1300 μm extending over $\sim 80' \times 68'$, Launhardt et al. (1996) have identified some of the dust emission peaks around NGC 2023 and NGC 2024 with the CS cores detected by Lada, Bally, & Stark (1991a). Of the existing FIR observations of NGC 2023 and NGC 2024, the IRAS observations in the first place trace relatively warmer dust (≥ 30 K) and secondly have angular resolutions varying with wavelength, so that calculating the temperature distribution from the observed flux densities is not straightforward. On the other hand, the 1300 μm observations, though of high angular resolution, trace extremely cold dust. This has motivated us to carry out simultaneous high angular

resolution far-infrared observations at 205 and 138 μm with identical beams so that the density and temperature distribution of dust emitting significantly in this intermediate regime can be reliably quantified.

The rest of the paper is organized as follows: Section 2 describes the observations and Section 3 contains the results and discussion of the same.

2. Observations

2.1. The 205 & 138 μm maps

Orion molecular clouds A and B (south) were observed using a two-band far-infrared photometer system (PHT-12) at the Cassegrain focus of the TIFR 1-meter (f/8) balloon-borne telescope. The far-infrared telescope was flown from the TIFR Balloon Facility, Hyderabad, in central India (Latitude = $17^\circ 47' \text{N}$, Longitude = $78^\circ 57' \text{E}$) in November, 1995. Details of the telescope and the observational procedure have been given by Ghosh et al. (1988). The photometer (PHT-12) consists of 12 composite silicon bolometers, each having a FOV of $1.6'$ and arranged in a 3×2 array for each band. These arrays are cooled to 0.3 K by liquid ^3He . The sky was chopped at 10 Hz by wobbling the secondary mirror. The chopper throw was $4.2'$ along the cross-elevation axis. The same region of the sky was viewed simultaneously in two bands. The effective wavelengths for the two bands for a λ^{-2} emissivity law and a temperature of 35 K are 205 μm and 138 μm . Saturn was observed for absolute flux calibration (Ghosh et al. 1988) as well as determination of the instrumental Point Spread Function (PSF) including the effect of sky chopping.

The simultaneous mapping in the two FIR bands was carried out by using raster scans in which the telescope was continuously moved along the cross-elevation axis and stepped in elevation. The chopped FIR signals were gridded in a two dimensional sky matrix (elevation $\times$ cross-elevation) with a cell size of $0.3' \times 0.3'$. The observed chopped signal matrix was deconvolved using an indigenously developed procedure based on the Maximum Entropy Method (MEM) similar to that of Gull & Daniell (1978) (see Ghosh et al. 1988, for details). The FWHM sizes of the deconvolved maps of the point-like source (Saturn) are found to be $1.6' \times 1.9'$ and $1.6' \times 1.8'$ in the 138 and 205 μm bands respectively. An optical photometer was used at the Cassegrain focal plane to improve the absolute positioning of the telescope during the flight. In addition for all the observations presented here,

the signal from this optical photometer measured during the far-infrared scans have been used during off-line analysis to further improve the aspect information. The absolute positional accuracy achieved in the FIR maps is ~ 0.5 .

2.2. The 57 μm map

Limited regions of both Orion A (OMC-1 & 2) and Orion B (NGC 2024) were also observed at 57 μm in an earlier flight (November, 1990) of the TIFR balloon-borne telescope, using a different photometer (PHT-2). This two band photometer, PHT-2, was based on two liquid ^4He cooled Ge(Ga) bolometers. The FIR passbands were 45–70 μm and 110–160 μm with effective wavelengths 57 μm and 150 μm for an input source spectrum defined by a greybody of temperature 35 K and λ^{-2} emissivity. A 2.4 (dia) region of the sky was simultaneously viewed in both the FIR bands. More details of PHT-2 can be found in Verma et al. (1994). Here, only the observations at 57 μm are presented, since the other band is very similar to the shorter waveband of PHT-12 which has better sensitivity. The PHT-2 was calibrated by observing Jupiter. Some preliminary results from observations of the Orion region have been presented by Das et al. (1993). The final results are presented here. The achieved angular resolution in the image processed maps is ~ 1.5 as determined from the observations of Jupiter.

2.3. The HIRES processed IRAS maps

To supplement our balloon-borne observations, we have used the IRAS survey data in all wavebands (12, 25, 60 and 100 μm) for the regions mapped by us in Orion B (south). These data were HIRES processed (Aumann, Fowler, & Melnyk, 1990) at the Infrared Processing and Analysis Center (IPAC¹, Caltech) for improving the angular resolutions of the raw maps.

3. Results & Discussion

3.1. Analysis of FIR data

Figure 1 and Figure 2 respectively show the regions (indicated by the dotted boundary) in Orion B (south) and Orion A mapped at 205 and 138 μm . Regions extending over $\sim 42' \times 38'$ in Orion A and $\sim 38' \times 53'$

in Orion B (south) have been observed at both these FIR wavelengths. The 57 μm maps for both sources cover relatively smaller regions. Reliable dust temperature $T(138/205)$ maps for selected regions have been generated utilising the simultaneity of observation and nearly identical beams at 205 and 138 μm . The dust emissivity law has been assumed to be $\epsilon_\nu \propto \nu^\beta$ and the method used is briefly described in Appendix 1. For most star forming regions the emissivity exponent at longer wavelengths ($> 100 \mu\text{m}$) is found to be between 1.5 to 2.5 and the calculated properties of most common types of dust grains (Draine & Lee 1984) show the same at these wavelengths. Based on these facts we have chosen $\beta = 2$ for deriving the $T(138/205)$ distribution. Considering the uncertainties in the measured flux densities and the wavelength regime probed, the temperature computed has an uncertainty of $\sim \pm 2\text{K}$ between 14 K and 50 K and of $\sim \pm 5\text{K}$ between 50 and 70 K. The intensity ratio of our FIR bands is rather insensitive to temperatures below 10 K and above 70 K. Above 70 K the estimated temperature has an uncertainty of $\sim \pm 8\text{K}$. The morphology of the spatial distribution of T_d is insensitive to the assumed value of β . Only the numerical values of T_d depend on β , e.g., a temperature of 15 K for $\beta = 2$ corresponds to 17 K for $\beta = 1$ and $T_d = 30$ K for $\beta = 2$ corresponds to 50 K for $\beta = 1$. Distribution of τ_{150} , the optical depth at 150 μm for Orion B has also been generated using these dust temperatures and $\beta = 2$.

We have adopted a well-defined extraction algorithm to identify discrete sources in these maps. This algorithm is based on identification of a local maximum followed by a critical study of the growth curve of the integrated flux densities in consecutive annular rings with the maximum at the center. The reliability of the sources detected in each map has been ensured by the following procedure. The entire observation for each region has been split into two independent datasets (by considering only the alternate scan lines in each raster) and processed (deconvolved) separately to generate two maps. The peaks detected in both these maps within a predetermined positional error have been used to quantify the dynamic range and also to validate the detected sources. The final maps presented combine all data and positions of the “confirmed sources” in the composite maps have been determined. These sources have signals more than 10 times the estimated noise level of the respective maps. In this paper the method of presenting the sources is as follows : Peaks identified in any two TIFR wave-

¹IPAC is funded by NASA as part of the IRAS extended mission program under contract to JPL.

bands are associated if they lie within $1'$. For association with IRAS (HIREs) sources the same parameter has been taken to be $1'.5$. The coordinates quoted for a particular source are for the position of the peak at the longest (TIFR) wavelength at which it is detected.

3.2. Orion A

Figure 2 shows the 205 and 138 μm intensity distribution of the mapped region in Orion A. The maps include the Orion Molecular Clouds 1, 2, 3 and the source IRAS 05327-0457. Figure 3 shows the intensity map for a limited region in Orion A at 57 μm . Table 1 presents the coordinates and flux densities (in a circle of $3'$ dia) for all the discrete sources detected in the mapped regions of Orion A at 205 and 138 μm . In the same table we also present the longer wavelength associations of these sources along with the measured flux densities at 1300 μm (Chini et al. 1997). For the discrete sources detected at 138, 205 and 1300 μm the dust temperature (T_d) and the emissivity exponent (β) have been determined by fitting a single temperature greybody spectrum. T_d and β for A8 have been determined in a similar manner by using the flux densities measured at 450, 790 and 1100 μm by Goldsmith, Bergin, & Lis (1997) along with those from the present work. For all other sources with flux densities measured only at 205 and 138 μm T_d has been determined assuming $\beta = 2$. For a source not detected in any one of the wavebands (marked by $\dagger$ in Table 1), we give the flux density obtained by integrating at the position of the source as detected in the other waveband(s). Table 2 presents the positions, flux densities and associations of these sources detected at 57 μm . The source IRAS 05327-0457 has been studied in detail in the far-infrared by Mookerjee et al. (2000); here we concentrate mainly on OMC-1, 2 and 3.

Figure 4 shows the temperature distribution as calculated from the observations at 205 and 138 μm for OMC-1 and 2. The temperature in OMC-1 is as high as ~ 70 K close to the Orion nebula. The temperature of the OMC-1 and the bar region as calculated by Werner et al. (1976) from the ratio of 50 and 100 μm flux densities is found to be between 55 and 85 K. For the same region, the temperature calculated from our 205 and 138 μm fluxes is between 45 and 70 K. This is consistent with the fact that the longer wavelength observations presented here are tracing colder dust component as compared to those traced by the 50 and 100 μm observations. The temperature drops to an average of ~ 25 K in OMC-2 and is as low as 15 K in OMC-3

(not shown in Figure). The temperature map has been generated by assuming the dust to be optically thin but explicit calculation of optical depth (at 150 μm) very close to the global intensity peak of OMC-1, shows the regions to be optically thick. In such cases, the temperatures quoted would be underestimates by as much as 5 K. However, for OMC-1, 2 and 3, the basic morphology of the isotherms remains unaltered.

Orion A is an ensemble of cores with a variety of temperatures, physical sizes and masses. This variety in physical conditions makes it conducive to harbouring stars in different stages of formation and evolution and hence makes it astrophysically interesting. The far-infrared maps presented here also support this variety by showing morphological characteristics strongly dependent on the dust temperature. While the 205 and 138 μm maps tracing colder dust detect FIR emission from OMC-2 and OMC-3 very clearly, the 57 μm map fails to do so.

OMC-1 (A8 in Table 1 and # 1 in Table 2) is the brightest source detected at all wavelengths. As mentioned earlier, the submillimeter fluxes at 450, 790 and 1100 μm (Goldsmith, Bergin, & Lis 1997) have been used along with the 57, 138 and 205 μm fluxes to determine T_d and β for this source. The dust temperature obtained by this method is 68 K and the β is 0.5. If instead we do a similar fitting omitting the flux density at 57 μm we get $T_d = 30$ K and $\beta = 1.1$. Thus the high temperature and low β as presented in Table 1 might be due to the effect of attempting to explain the emission over a wide range of wavelengths using a single temperature greybody spectrum. In reality there is probably a cold diffuse component of dust as well which contributes to the emission at longer wavelengths. In the OMC-1 region the maps at 205 & 138 μm also detect the source A5. Although the position of this source at 205 and 138 μm differ by $\sim 1'.5$, we have associated the two, since there is no other nearby source of similar strength. We also note that the temperature of the source A5 is ~ 80 K, which is even higher than the temperature of A8.

The peaks # 2 and # 3 detected at 57 μm (see Table 2) are not detected at the longer wavelengths. The source # 2 is associated with the HII region M43 and the intensity distribution around it is similar to that observed at 60 μm by Smith, Harper, & Loewenstein (1987). The source # 3 is due to the heating of the dust in the photodissociation region around the trapezium stars. Northern tip of the emission detected at 57 μm is clearly displaced to the east with respect to the inte-

gral shaped emission ridge detected in OMC-2 and 3. We can discern local enhancements in T(138/205) map close to the positions of the peaks #2 and #3. These strongly indicate the existence of hotter dust towards the east.

Of the observations presented here OMC-2 and 3 are detected only at the longer wavelengths (205 and 138 μm). In OMC-2 we detect two sources (A10 and A11) at 138 μm and only one source (A12) at 205 μm . As suggested by Table 1 the sources detected at our resolution are each associated with more than one sub-millimeter (1300 μm) sources. We note that whereas A10 and A11 are associated with two mutually exclusive groups of submillimeter sources, A12 is associated with sources from both groups. The fact that we detect A10 and A11 only at 138 μm and A12 only at 205 μm could be due to the presence of a diffuse cold component which reduces the contrast between the sources and the intermediate region. This could effectively smoothen out the smaller peaks (like A10 and A11) in this region, thus resulting in only one peak (A12) at 205 μm . The flux densities at 1300 μm of the detected sub-mm sources have been calculated using the peak fluxes and size of the individual sources (refer to Table 1 of Chini et al. 1997). For each FIR source the contribution at 1300 μm from all associated sources are combined and presented in Table 1. For all sources detected in OMC-2 a single temperature greybody spectral fitting gives $20\text{K} < T_d < 25\text{K}$ and $1.6 < \beta < 2.0$. Both these are consistent with the results of Chini et al. (1997). Recent radio observations at 3.6 cm by Reipurth, Rodriguez, & Chini (1999) have detected sources coinciding with the peaks in dust emission. These associations are also presented in Table 1. The radio emission further substantiates that the sources detected in OMC-2 though reasonably cold are not only regions of high density but also have YSOs embedded in them.

We have detected one source (A9) in the OMC-3 region. Positionally it is coincident with the sub-mm source MMS 6 (Chini et al. 1997), the latter being the brightest of all the sub-mm sources detected in OMC-3. The flux density at 1300 μm for the source A9 is estimated in the same manner as above. A greybody spectrum fitted to the measured flux densities gives $T_d = 15\text{ K}$ and $\beta = 2.0$. A9 is thus the coldest source in Orion A, both T_d and β estimated here are consistent with Chini et al. (1997) and Lis et al. (1998).

The total far-infrared luminosity has been estimated using the temperature map (Fig. 4) for the region with

OMC-1 and OMC-2 and the intensity maps of the same region. The dust emissivity exponent assumed for this calculation is 2.0. From the FIR observations presented here the total luminosity detected in OMC-1 and 2 is $\sim 3.5 \times 10^5 L_\odot$. For a similar region the luminosity based on 60 and 100 μm observations as given by Stacey et al. (1993) is $\sim 5 \times 10^5 L_\odot$. The diffuse emission at these wavelengths has been calculated by subtracting the contribution of the discrete sources and is about 25% of the total luminosity.

The molecular cloud OMC-1 has long been the most extensively studied region in Orion A. OMC-2 and 3 have been studied only in the recent past and have been found to be rich and interesting as a star forming region. Our observations provide useful constraints on the intensity and temperature distribution of this region.

3.3. Orion B (south)

3.3.1. Region around NGC 2024

Figure 5 presents intensity maps of the region around NGC 2024 at 205, 138, 100 and 57 μm . The map at 57 μm is for a smaller region, but brings out the main features of NGC 2024 at this wavelength. Table 3 presents the measured flux densities at the 4 IRAS and 3 TIFR wavelengths for the sources detected in this region. The sub-mm and IRAS Point Sources associated with these FIR sources, along with the sub-mm flux densities measured at 1300 μm by Launhardt et al. (1996) are presented in the Table 3. In all 7 sources have been identified with sources in the IRAS Point Source Catalog (PSC) and 2 sources (B3 and B9) have been found to have sub-millimeter (1300 μm) counterparts. For sources not detected in one or two of the TIFR bands (marked by † in Table 3) we have presented the flux densities obtained by integrating in a circle around the position of the source as detected at the other wavelength(s). We have fitted a single temperature greybody spectrum for all sources for which measurement of flux densities at 3 or more wavelengths longer than 50 μm exist. The resulting T_d and β are also presented in the same table. For the remaining sources T_d is determined using $\beta = 2$. We find that the emissivity exponent (β) for some sources (B6, B10 and B12) are very small (≤ 0.5). In addition, for B6 and B12 the fitted T_d is also very high. This indicates that for these sources possibly a two or multi-temperature greybody spectrum is more appropriate.

Observations at $1300\ \mu\text{m}$ by Mezger et al. (1988) have revealed 6 protostellar condensations within an area of $3' \times 3'$ around NGC 2024. The far-infrared maps presented here being of limited angular resolution do not resolve these condensations. However the $205\ \mu\text{m}$ and also the $100\ \mu\text{m}$ maps show evidence of north-south extension with the northern tip having a slight westward bend (position of FIR-1, 2 ; Mezger et al. 1988). Positionally B9 coincides with FIR-5 (brightest condensation detected by Mezger et al. 1988) and also with LBS 33SM, a CS core detected by Lada, Bally, & Stark (1991a) (hereafter LBS), later detected at $1300\ \mu\text{m}$ by Launhardt et al. (1996). The HIRES map at $100\ \mu\text{m}$ shows possible signs of anomaly, particularly near the peak. The flux density measured (in $3'$ dia) from the $100\ \mu\text{m}$ map shows a very low value of $\sim 18\ \text{kJy}$ compared to both the IRAS PSC value ($35.3\ \text{kJy}$) and results from the KAO observations by Thronson et al. (1984) ($85\ \text{kJy}$). We have used the measurements of Thronson et al. (1984) in the Table 3 as well as for determining T_d and β for B9.

Figure 6 shows the dust temperature distribution for this region. It has been generated using the same method as for Orion A. It shows several isolated peaks with temperatures varying between 60 to 70 K for most of them. The source B3 is quite cold with a maximum of 30 K and going below 15 K towards the edges. The τ_{150} map (Figure 6) shows very interesting structure close to the main source B9. It distinctly shows that the region around B9 has an elongated structure with indications of at least 2 condensations. This structure is very similar to the structure shown by the NH_3 column density map generated by Schulz et al. (1991). This very well substantiates the fact that these are indeed protostellar condensations since they are so luminous in the infrared as well. The cold source B3 is found to be mostly optically thin (peak 0.07) at $150\ \mu\text{m}$.

The total luminosity of the region calculated using the same method as for Orion A (Section 3.1) is $1.0 \times 10^5 L_\odot$. The diffuse emission contributes about 50% (at both the wavelengths) of the total luminosity. This is substantially larger than the value measured in Orion A (Sec 3.1) and in other young Galactic star forming regions ($\sim 35\%$ in W31; Ghosh et al. 1989). The morphology of the [CII] ($158\ \mu\text{m}$) emitting region in Orion B (south) as observed by Jaffe et al. (1994) agrees very well with the maps presented in this paper. The total [CII] luminosity detected for their assumed distance of 415 pc is $230 L_\odot$, while we have estimated the total far-infrared luminosity for a distance of 450

pc. Correcting for this difference in distances assumed we find the ratio of the [CII] luminosity to the total FIR luminosity to be 3×10^{-3} .

The observations presented here, being on a large scale and of good angular resolution would be useful as guidelines in understanding the star formation scenario vis-a-vis the temperature distribution for future large scale sub-millimeter continuum observations.

3.3.2. Region around NGC 2023

Figure 7 shows intensity maps of the region surrounding NGC 2023 at 205 and $138\ \mu\text{m}$ alongwith HIRES maps at 100 and $60\ \mu\text{m}$. The maps at 205 and $138\ \mu\text{m}$ are generated by masking NGC 2024 prior to MEM deconvolution of the larger map so that details of the fainter source NGC 2023 are better visible and comparison with 100 and $60\ \mu\text{m}$ HIRES maps is easier. The source NGC 2023 appears to have a double structure at $205\ \mu\text{m}$. It is of interest to note that the map of fluorescent H_2 observed by Gatley et al. (1987) exhibits a shell structure with main peaks separated by similar extent as B13 and B14. The intensity map at $138\ \mu\text{m}$ however shows only a single source close to the brighter source (B14) at $205\ \mu\text{m}$. Diffuse emission extending southward from the main source is clearly visible at 60 , 100 and $205\ \mu\text{m}$, while at $138\ \mu\text{m}$ the limited dynamic range of the map restricts this emission to a seemingly isolated peak. Table 3 presents the positions, flux densities and associated sources for the 2 peaks (B13 and B14) detected in this region. The coordinates of the global peaks in the 60 and $100\ \mu\text{m}$ maps is found to be $\sim 5^{\text{h}} 39^{\text{m}} 7'' - 2^{\circ} 17' 44''$. These global peaks are separated from B14 by more than $1/5$. Since there is no other detected source of such brightness in the neighbourhood and observations at 100 and $138\ \mu\text{m}$ trace dust of similar temperature we have associated B14 with NGC 2023. Both sources B13 and B14 have been detected at $1300\ \mu\text{m}$ by Launhardt et al. (1996), but there is no IRAS detection of B13. For both sources we have used the available flux densities at wavelengths longer than $60\ \mu\text{m}$ to estimate T_d and β from a single temperature greybody spectrum (Table 3). The flatness of β for B14 indicates the existence of dust components of at least two temperature ranges. The source B13 is found to be reasonably cold ($T_d = 16\ \text{K}$), thus justifying it's non-detection in the IRAS wavebands.

Figure 8 shows the $T(138/205)$ and τ_{150} maps for the region around NGC 2023 (derived assuming $\beta =$

2). The T(138/205) map shows a peak which does not coincide with the intensity peaks at 205 and 138 μm . The T(138/205) peak, however coincides with the peak positions at 60 and 100 μm . The temperature of most of the region is around 30 K and in places goes down to values as low as ~ 20 K. The τ_{150} map more or less traces the high intensity regions. The τ_{150} displayed for regions which lie outside the contours of T(138/205) map are for an assumed temperature of 10 K (not plotted in the T(138/205) map).

NGC 2023, being a well known reflection nebula with a photodissociation region (excited by HD 37903) has been studied in various molecular and fine structure ([CII] at 158 μm) lines. The continuum maps presented in Figure 7 are morphologically similar to the C¹⁸O map by Wyrowski et al. (1997) and CO (J=2-1) map by White et al. (1990). The [CII] observations at 158 μm by Howe et al. (1991) and Jaffe et al. (1994) detect emission which is approximately 25 % of the emission from NGC 2024. The rise in temperature, T(138/205), to the north-west of the main peak is explained by the presence of the exciting star HD 37903. The total luminosity measured in this region is $1.5 \times 10^3 L_{\odot}$. The diffuse emission contributes about 35% of the total emission, which is smaller than the contribution of diffuse emission around NGC 2024. The total [CII] luminosity from the region as detected by Howe et al. (1991) is $7 L_{\odot}$ and is equal to $\sim 0.5\%$ of the total FIR luminosity estimated here. This is consistent with results from most of the Galactic star forming regions as given by Howe et al. (1991). Detection of low temperature in the region around NGC 2023 alongwith emission in the far-infrared, support ongoing formation of low to intermediate stars in the region. Indication of more than 50 % of the emission being in the condensations further indicates star formation in clumps in this region.

4. Summary

In this paper we have presented large scale high angular resolution ($\sim 1'$) far-infrared (205 and 138 μm) maps of the Orion A and Orion B (south) molecular clouds. For limited regions, maps at 57 μm have also been presented. HIREs processed IRAS images at 100 and 60 μm of the Orion B have been used for comparison. The regions covered include OMC-1, 2, 3 in Orion A and NGC 2024 and 2023 in Orion B (south). In all 15 condensations in Orion A and 14 condensations in Orion B (south) have been identified. In OMC-

3 the very cold source MMS 6 (Chini et al. 1997) was detected at both 205 and 138 μm and the temperature was determined to be $\sim 15 \pm 2$ K. The far-infrared counterparts of the extremely dense CS cores (Lada, Bally, & Stark 1991a) LBS 33, LBS 34, LBS 36 and LBS 40 were detected in Orion B (south). The total far-infrared luminosity and the relative contribution of diffuse emission to the luminosity of these regions have also been quantified.

We thank B. Das for his help during initial stages of analysis of the 57 μm data and S.L. D'Costa, M.V. Naik, D.M. Patkar, M.B. Naik, S.A. Chalke, G.S. Meshram & C.B. Bakalkar for their support for the experiment. The members of TIFR Balloon Facility (Balloon group and Control & Instrumentation group), Hyderabad, are thanked for their roles in conducting the balloon flights. Thanks are due to IPAC for providing HIREs processed IRAS data. We thank the anonymous referee for the suggestions which have improved the quality of the paper.

A. Estimation of dust temperature distribution using the two band FIR observations

As mentioned in Section 2.1 the spatial intensity distribution in each band is gridded with pixels of size 0.3×0.3 . Only those pixels which have signals exceeding 5 times the noise level of the corresponding map are considered to be “valid” for the purpose of finding the dust temperature distribution. Each intensity map is first smoothed by taking a running average over 3×3 “valid” pixels. This is done to be extremely conservative about the derived spatial structures in the temperature maps. Since the two intensity maps refer to identical portions of the sky and the beams are also almost identical, a “pixel by pixel” ratio between the two maps gives reliable dust temperatures.

For optically thin emission the observed flux density at any wavelength can be written as

$$F_\nu = \Omega B_\nu(T_d) \tau_\nu \quad (\text{A1})$$

where Ω is the solid angle of the region under consideration.

We assume $\tau_\nu \propto \nu^\beta$. Since the pixel sizes are identical the ratio of flux densities at any two wavelengths would be:

$$\frac{F_{\nu_1}}{F_{\nu_2}} = \left(\frac{\nu_1}{\nu_2} \right)^{3+\beta} \frac{[\exp(h\nu_2/kT_d) - 1]}{[\exp(h\nu_1/kT_d) - 1]} \quad (\text{A2})$$

Thus for an assumed emissivity exponent (β) the ratio of the flux densities at any two wavelengths is a function of the dust temperature only. The temperature is computed using a lookup table containing the calculated ratios of flux densities for various temperatures and an assumed β . This process is repeated for all pixels to generate the dust temperature distribution map.

REFERENCES

- Aumann, H. H., Fowler, J. W., & Melnyk, M. 1990, *AJ*, 99, 1674
- Castets, A., & Langer, W. D., 1995, *A&A*, 294, 835
- Chini, R., Reipurth, B., Ward-Thompson, D., Bally, J., Nyman, L. -A., Sievers, A., & Billawala, Y. 1997, *ApJ*, 474, L135
- Das, B., Ghosh, S. K., Rengarajan, T. N., Verma, R. P., Iyengar, K. V. K., & Tandon, S. N. 1993, *Bull. Astron. Soc. India*, 21, 587
- Draine, B. T., & Lee, H. M. 1984, *ApJ*, 285, 89
- Gatley, I., Hasegawa, T., Suzuki, H., Garden, R., Brand, P., Lightfoot, J., Glencross, W. Okuda, H., & Nagata, T. 1987, *ApJ*, 318, L73
- Ghosh, S. K., Iyengar, K. V. K., Rengarajan, T. N., Tandon, S. N., Verma, R. P., & Daniel, R. R. 1988, *ApJ*, 330, 928
- Ghosh, S. K., Iyengar, K. V. K., Rengarajan, T. N., Tandon, S. N., Verma, R. P., Daniel, R. R., & Ho, P. T. P. 1989, *ApJ*, 347, 338
- Goldsmith, P. F., Bergin, E. A., & Lis, D. C. 1997, *ApJ*, 491, 615
- Gull, S. F., & Daniell, G. J. 1978, *Nature*, 272, 686
- Howe, J. E., Jaffe, D. T., Genzel, R., & Stacey, G. J. 1991, *ApJ*, 373, 158
- Jaffe, D. T., Zhou, S., Howe, J. E., Herrmann, F., Madden, S. C., Poglitsch, A., van der Werf, P. P., & Stacey, G. J. 1994, *ApJ*, 436, 203
- Johnstone, D., & Bally, J. 1999, *ApJ*, 510, L49
- Lada, E. A., Bally, J., & Stark, A. A. 1991a, *ApJ*, 368, 432.
- Lada, E. A., DePoy, D. L., Evans, N. J., & Gatley, I. 1991b, *ApJ*, 371, 171.
- Launhardt, R., Mezger, P. G., Haslam, C. G. T., Kreysa, E., Lemke, R., Sievers, A., & Zylka, R. 1996, *A&A*, 312, 569
- Lis, D. C., Serabyn, E., Keene, J., Dowell, C. D., Benford, D. J., Phillips, T. G., Hunter, T. R., & Wang, N. 1998, *ApJ*, 509, 299
- Mezger, P. G., Chini, R., Kreysa, E., Wink, J. E., & Salter, C. J. 1988, *A&A*, 191, 44
- Mezger, P. G., Sievers, A. W., Haslam, C. G. T., Kreysa, E., Lemke, R., Mauersberger, R., & Wilson, T. L. 1992, *A&A*, 256, 631
- Mookerjee, B., Ghosh, S. K., Rengarajan, T. N., Tandon, S. N., & Verma, R. P. 2000, *ApJ*(in press)
- Reipurth, B., Rodriguez, L. F., & Chini, R. 1999, *AJ*, 118, 983
- Schulz, A., Gusten, A., Zylka, R., & Serabyn, E. 1991, *A&A*, 246, 570
- Smith, J., Harper, D. A., & Loewenstein, R. F. 1987, *ApJ*, 314, 76
- Stacey, G. J., Jaffe, D. T., Geis, N., Genzel, R., Harris, A. I., Poglitsch, A., Stutzki, J., & Townes, C. H. 1993, *ApJ*, 404, 219
- Thronson, H. A., Lada, C. J., Schwartz, P. R., Smith, H. A., Smith, J., Glaccum, W., Harper, D. A., & Loewenstein, R. F. 1984, *ApJ*, 280, 154
- Thronson, H. A., Harper, D. A., Bally, J., Dragan, M., Mozurkewich, D., Greenhouse, M. A., Schwartz, P. R., Smith, H. A., Bieging, J. H., Loewenstein, R. F., & Lada, C. J. 1986, *AJ*, 91, 1350
- Verma, R. P., Bisht, R. S., Ghosh, S. K., Iyengar, K. V. K., Rengarajan, T. N., & Tandon, S. N. 1994, *A&A*, 284, 936
- Werner, M. W., Gatley, I., Harper, D. A., Becklin, E. E., Loewenstein, R. F., Telesco, C. M., & Thronson, H. A. 1976, *ApJ*, 204, 420
- White, G. J., Sanderson, C., Monteiro, T. S., Richardson, K. J., & Hayashi, S. S. 1990, *A&A*, 227, 200
- Wyrowski, F., Walmsley, C. M., Natta, A., & Tielens, A. G. G. M. 1997, *A&A*, 324, 1135

Figure Captions

Contour levels are same as in Fig. 7c.

Fig. 1.— Intensity map of Orion B (South) at 205 μm . Dotted boundary shows the region mapped. The peak is 2.50 kJy/sq. arcmin. The contour levels are at 90 to 30% (in steps of 20), and 20, 10, 5, 2.5, 1.25 and 0.63% of the peak.

Fig. 2.— The intensity maps of Orion A. Peaks are 9.00 kJy/sq. arcmin at 205 μm and 29.1 kJy/sq. arcmin at 138 μm . The contour levels are same as in Figure 1, with an additional contour at 0.31% of the peak. The lowest contour level is 3 times the measured noise.

Fig. 3.— The intensity map of the Orion A at 57 μm . The peak is 59.6 kJy/sq. arcmin. The contour levels are same as in Figure 2.

Fig. 4.— The dust temperature map for the region covering OMC-1 and 2, assuming a dust emissivity of $\epsilon_\nu \propto \nu^2$. The lowest temperature found in this region is 14 K. The numbers written on the contours refer to the temperatures, with the highest contour displayed being at 70 K.

Fig. 5.— The intensity maps of the region around NGC 2024. Peaks are (a) 2.50 kJy/sq. arcmin at 205 μm , (b) 7.53 kJy/sq. arcmin at 138 μm , (c) 3.80 kJy/sq. arcmin at 100 μm and (d) 19.1 kJy/sq. arcmin at 57 μm . The contour levels (a) and (b) are same as in Figure 1. Levels in (c) are 60 to 10 % (in steps of 10) and 5, 2.5 and 1% of its peak. Levels in (d) are 90, 75, 60, 40, 20, 10, 7.5, 5, 3.5, 2, 1 and 0.7% of the peak.

Fig. 6.— (a) Same as in Fig 4, except being for the region around NGC 2024. (b) τ_{150} map for the same region. Contour levels are 90 to 10 % (in steps of 10) and 5, 2.5 and 1% of its peak (0.24).

Fig. 7.— The intensity maps for the region around NGC 2023. The peaks are (a) 325 Jy/sq. arcmin at 205 μm , (b) 329 Jy/sq. arcmin at 138 μm , (c) 511 Jy/sq. arcmin at 100 μm and (d) 786 Jy/sq. arcmin at 60 μm . The contour levels in (a) are 95%, 90 to 10% (in steps of 10) and 5% of the peak and in (b) are 95%, 90 to 30 % (in steps of 10) of the peak. The contour levels in (c) and (d) are the same as in Fig. 6b except for an additional contour at 95 % of the peak.

Fig. 8.— (a) T(138/205) map for the region around NGC 2023. The highest contour displayed is at 40 K. (b) τ_{150} map for NGC 2023 with peak value of 0.14.

TABLE 1
POSITION, FLUX DENSITIES (IN 3' DIA), ASSOCIATIONS, T_d & β OF SOURCES IDENTIFIED IN ORION A AT 205
AND 138 μM

Source #	Position ^a (1950)		Association ^b	F_{138} kJy	F_{205} kJy	F_{1300} Jy	T_d^c K	β^c
	R.A.	Dec.						
A1	5 31 51.6	- 5 06 22	IRAS 05318-0506	1.26	0.58	...	29	...
A2	5 32 02.0	- 4 56 30	...	1.24 [†]	0.96	...	20	...
A3	5 32 04.6	- 5 19 14	...	0.83	0.39	...	28	...
A4	5 32 11.6	- 5 02 28	...	0.56	0.36	...	22	...
A5	5 32 32.8	- 5 22 57	...	2.65	0.68	...	80	...
A6	5 32 43.7	- 4 57 42	IRAS 05327-0457	2.12	0.96	...	30	...
A7	5 32 47.3	- 4 58 33	...	2.06	0.86 [†]	...	>32	...
A8	5 32 47.7	- 5 24 51	OMC-1	112.50	35.70	...	68 ^d	0.5 ^d
A9	5 32 55.6	- 5 04 00	OMC-3, MMS 4,5,6, VLA 3	0.42	0.53	4.77	15	2.0
A10	5 32 59.5	- 5 14 25	FIR 6a,6b,6c,6d, VLA 14	2.22	1.44 [†]	7.45	25	1.6
A11	5 33 01.4	- 5 11 16	FIR 2,3,4, VLA 11,12	1.88	1.50 [†]	6.94	20	1.9
A12	5 33 01.6	- 5 13 01	FIR 4,5,6a,6b	1.98 [†]	1.71	10.4	20	1.8
A13	5 33 07.1	- 5 01 49	...	0.43 [†]	0.54	...	15	...

[†]Not detected as a local peak in this waveband. Flux density is obtained by integrating in a circle of 3' dia around the coordinates tabulated.

^aUnits of right ascension are hours, minutes and seconds, and units of declination are degrees, arcminutes, and arcseconds.

^bIRAS : IRAS PSC ; FIR & MMS : Chini et al. (1997); VLA : Reipurth, Rodriguez, & Chini (1999).

^cSingle temperature greybody spectrum; if flux densities at only 2 bands are available emissivity $\epsilon_\nu \propto \nu^2$ is assumed.

^dUsing F_{57} , F_{138} & F_{205} (this work) and $F_{450} = 19.8$ kJy, $F_{790} = 2.671$ kJy & $F_{1100} = 0.989$ kJy (Goldsmith, Bergin, & Lis 1997).

TABLE 2
POSITION AND FLUX DENSITIES (IN 3' DIA) OF SOURCES IDENTIFIED IN ORION A AT 57 μM

Source #	Position ^a (1950)		F_{57} kJy	Associations
	R.A.	Dec.		
1	5 32 47.3	- 5 24 27	219	OMC-1
2	5 33 00.2	- 5 16 18	5.91	M43
3	5 33 03.3	- 5 21 46	3.99	...

^aUnits of right ascension are hours, minutes and seconds, and units of declination are degrees, arcminutes, and arcseconds.

TABLE 3
POSITION, FLUX DENSITIES (IN $3'$ DIA), ASSOCIATIONS, T_d & β OF SOURCES IDENTIFIED IN ORION B AT 205
AND $138 \mu\text{M}$

Source #	Position ^a (1950)		Association ^b	F ₁₂ Jy	F ₂₅ Jy	F ₅₇ kJy	F ₆₀ kJy	F ₁₀₀ kJy	F ₁₃₈ kJy	F ₂₀₅ kJy	F ₁₃₀₀ Jy	T _d ^c	β ^c
	R.A.	Dec.											
NGC 2024 region													
B1	5 38 21.4	- 1 50 19	05383-0150	60	65	NM ^d	0.95	1.40	0.67	0.29	...	32	2.5
B2	5 38 21.9	- 1 54 24	...	68	68	NM	1.17	...	1.04	0.30	...	25	3.8
B3	5 38 43.0	- 1 49 55	05387-0149	80	133	...	...	...	1.69	1.13	2.6	20	2.3
			LBS 40 SM										
B4	5 38 44.5	- 1 58 44	...	87	120	4.50 [†]	...	2.79	2.18	0.67	...	21	3.8
B5	5 38 52.6	- 1 46 10	05388-0147	89	...	NM	...	...	0.57 [†]	0.49	...	20	...
B6	5 38 57.2	- 1 54 00	...	110	...	5.51	...	...	1.89	0.90	...	95	0.4
B7	5 38 59.4	- 1 47 52	...	81	175	NM	2.83	3.59	1.25	0.37 [†]	...	27	3.9
B8	5 39 08.4	- 1 52 13	05391-0152	248	1060	6.62	10.74	...	4.24	1.31	...	32	2.8
B9	5 39 13.7	- 1 56 52	05393-0156	86	916	71.70	...	85.00 ^e	32.30	12.06	115	59	1.0
			LBS 33 SM										
B10	5 39 21.8	- 1 45 47	05392-0145	17	31	NM	0.32	...	0.83	0.59	...	35	0.5
B11	5 39 21.8	- 1 49 13	05393-0150	80	176	0.71 [†]	2.10	2.96	1.56	0.65	...	32	2.4
B12	5 39 28.5	- 2 01 36	...	...	...	...	...	0.77	0.51	0.27 [†]	...	80	0.2
NGC 2023 region													
B13	5 39 02.9	- 2 20 16	LBS 36 SM	25	52	NM	...	...	0.79 [†]	0.82	4.7	16	2.1
B14	5 39 14.6	- 2 19 08	LBS 34 SM	14	...	NM	...	...	1.93	1.41	12.5	25	1.3

[†]Not detected as a local peak in this waveband. Flux density is obtained by integrating in a circle of $3'$ dia around the coordinates tabulated.

^aUnits of right ascension are hours, minutes and seconds, and units of declination are degrees, arcminutes, and arcseconds.

^bLBS SM : CS cores (Lada, Bally, & Stark 1991a) also detected at $1300 \mu\text{m}$ by Launhardt et al. (1996). IRAS PSC for all other entries.

^cSingle temperature greybody spectrum; if fluxes at only 2 bands are available emissivity $\epsilon_\nu \propto \nu^2$ is assumed.

^dNot mapped at $57 \mu\text{m}$.

^eThronson et al. (1984).















